

ON GENERALIZED THE CONFORMABLE FRACTIONAL CALCULUS

M. Z. SARIKAYA¹, H. BUDAK¹, F. USTA^{1, §}

ABSTRACT. In this paper, we generalize the conformable fractional derivative and integral and obtain several results such as the product rule, quotient rule, chain rule.

Keywords: Conformable fractional derivative, conformable fractional integrals.

AMS Subject Classification: 23A33, 26A42

1. INTRODUCTION

An important point is that the fractional derivative at a point x is a local property only when α is an integer; in non-integer cases we cannot say that the fractional derivative at x of a function f depends only on values of f very near x , in the way that integer-power derivatives certainly do. Therefore it is expected that the theory involves some sort of boundary conditions, involving information on the function further out. To use a metaphor, the fractional derivative requires some peripheral vision. As far as the existence of such a theory is concerned, the foundations of the subject were laid by Liouville in a paper from 1832. The fractional derivative of a function to order α is often now defined by means of the Fourier or Mellin integral transforms. Various types of fractional derivatives were introduced: Riemann-Liouville, Caputo, Hadamard, Erdelyi-Kober, Grunwald-Letnikov, Marchaud and Riesz are just a few to name [7]-[10]. Recently a new local, limit-based definition of a so-called conformable derivative has been formulated in [1], [6] as follows

$$D_{\alpha}(f)(t) = \lim_{\varepsilon \rightarrow 0} \frac{f(t + \varepsilon t^{1-\alpha}) - f(t)}{\varepsilon}$$

proved the limits exists. Note that if f is fully differentiable at t , then the derivative is $D_{\alpha}(f)(t) = t^{1-\alpha} f'(t)$. The reader interested on the subject of conformable calculus is referred to the [1]-[7].

In this paper, we introduce a new fractional derivative which is generalized the results obtained in [1] and [6]. Then, we establish some basic tools for fractional differentiation and fractional integration. Furthermore, if $\alpha = 1$, the definition is equivalent to the

¹ Department of Mathematics, Faculty of Science and Arts, Düzce University, Düzce-TURKEY.
 sarikayamz@gmail.com, ORCID: <http://orcid.org/0000-0002-6165-9242>.
 hsyn.budak@gmail.com; ORCID:<http://orcid.org/0000-0001-8843-955X>.
 fuatusta@duzce.edu.tr; ORCID:<https://orcid.org/0000-0002-7750-6910>.

§ Manuscript received: August 25, 2017; accepted: December 21, 2017.

TWMS Journal of Applied and Engineering Mathematics, Vol.9, No.4 © Işık University, Department of Mathematics, 2019; all rights reserved.

classical definition of the first order derivative of the function f . Furthermore, it is noted that there are (α, a) -differentiable functions which are not differentiable.

2. DEFINITIONS AND PROPERTIES OF a -CONFORMABLE FRACTIONAL DERIVATIVE

In this section, we give a new definition and obtain several results such as the product rule, quotient rule and chain rule. We start with the following definition which is a generalization of the conformable fractional derivative.

Definition 2.1 (a -Conformable fractional derivative). *Given a function $f : [a, b] \rightarrow \mathbb{R}$ with $0 \leq a < b$. Then the “ a -conformable fractional derivative” of f of order α is defined by*

$$D_{\alpha}^a(f)(t) = \lim_{\varepsilon \rightarrow 0} \frac{f(t + \varepsilon t^{-\alpha}(t-a)) - f(t)}{\varepsilon(1 - at^{-\alpha})}, \quad (1)$$

for all $t > a$, $t^{\alpha} \neq a$, $\alpha \in (0, 1)$. If f is (α, a) -differentiable in some (a, b) , $a > 0$, $\lim_{t \rightarrow a^+} D_{\alpha}^a(f)(t)$ exist, then define

$$D_{\alpha}^a(f)(a) = \lim_{t \rightarrow a^+} D_{\alpha}^a(f)(t). \quad (2)$$

If the a -conformable fractional derivative of f of order α exists, then we simply say f is (α, a) -differentiable.

Theorem 2.1. *Let $0 \leq a < b$ and $\alpha \in (0, 1]$. If a function $f : [a, b] \rightarrow \mathbb{R}$ with is (α, a) -differentiable at $t_0 > a$, $t_0^{\alpha} \neq a$, then f is continuous at t_0 .*

Proof. Since $f(t_0 + \varepsilon t_0^{-\alpha}(t_0 - a)) - f(t_0) = \frac{f(t_0 + \varepsilon t_0^{-\alpha}(t_0 - a)) - f(t_0)}{\varepsilon(1 - at_0^{-\alpha})} \varepsilon(1 - at_0^{-\alpha})$, we have

$$\lim_{\varepsilon \rightarrow 0} f(t_0 + \varepsilon t_0^{-\alpha}(t_0 - a)) - f(t_0) = \lim_{\varepsilon \rightarrow 0} \frac{f(t_0 + \varepsilon t_0^{-\alpha}(t_0 - a)) - f(t_0)}{\varepsilon(1 - at_0^{-\alpha})} \lim_{\varepsilon \rightarrow 0} \varepsilon(1 - at_0^{-\alpha}).$$

Let $h = \varepsilon t_0^{-\alpha}(t_0 - a)$. Then we get

$$\lim_{\varepsilon \rightarrow 0} f(t_0 + h) - f(t_0) = D_{\alpha}^a(f)(t_0) \cdot 0,$$

which implies that f is continuous at t_0 . This completes the proof. \square

Having these definitions in hand we can present the following properties for (α, a) -differentiable functions:

Theorem 2.2. *Let $\alpha \in (0, 1]$ and f, g be (α, a) -differentiable at a point $t > a$, $t^{\alpha} \neq a$. Then*

- i. $D_{\alpha}^a(\lambda_1 f + \lambda_2 g) = \lambda_1 D_{\alpha}^a(f) + \lambda_2 D_{\alpha}^a(g)$, for all $\lambda_1, \lambda_2 \in \mathbb{R}$,
- ii. $D_{\alpha}^a(t^n) = \frac{nt^{n-1}(t-a)}{(t^{\alpha}-a)}$ for all $n \in \mathbb{R}$
- iii. $D_{\alpha}^a(c) = 0$, for all constant functions $f(t) = c$,
- iv. $D_{\alpha}^a(fg) = gD_{\alpha}^a(f) + fD_{\alpha}^a(g)$,
- v. $D_{\alpha}^a\left(\frac{f}{g}\right) = \frac{fD_{\alpha}^a(g) - gD_{\alpha}^a(f)}{g^2}$,
- vi. $D_{\alpha}^a(f \circ g) = f'(g(t))D_{\alpha}^a(g)(t)$ for f differentiable at $g(t)$,
- vii. If, in addition, f is a differentiable, then $D_{\alpha}^a(f)(t) = \frac{(t-a)}{(t^{\alpha}-a)}f'(t)$.

Proof. Parts *i* and *iii* follow directly from the definition. Now, we will prove *ii*, *iv*, *vi* and *vii*. For fixed and $t > a$, $t^\alpha \neq a$, we have

$$\begin{aligned} D_\alpha^a(f)(t) &= \lim_{\varepsilon \rightarrow 0} \frac{(t + \varepsilon t^{-\alpha}(t-a))^n - t^n}{\varepsilon(1 - at^{-\alpha})} \\ &= \lim_{\varepsilon \rightarrow 0} \frac{t^n + n\varepsilon t^{-\alpha}(t-a)t^{n-1} + O(\varepsilon^2) - t^n}{\varepsilon(1 - at^{-\alpha})} \\ &= \frac{nt^{n-1}(t-a)}{(t^\alpha - a)}. \end{aligned}$$

This completes the proof *ii*. Then, we will prove *iv*. For this purpose, since f, g are (α, a) -differentiable at a point $t > a$, $t^\alpha \neq a$, we obtain

$$\begin{aligned} D_\alpha^a(fg)(t) &= \lim_{\varepsilon \rightarrow 0} \frac{f(t + \varepsilon t^{-\alpha}(t-a))g(t + \varepsilon t^{-\alpha}(t-a)) - f(t)g(t)}{\varepsilon(1 - at^{-\alpha})} \\ &= \lim_{\varepsilon \rightarrow 0} \left[\frac{f(t + \varepsilon t^{-\alpha}(t-a)) - f(t)}{\varepsilon(1 - at^{-\alpha})} g(t + \varepsilon t^{-\alpha}(t-a)) \right] \\ &\quad + f(t) \lim_{\varepsilon \rightarrow 0} \frac{g(t + \varepsilon t^{-\alpha}(t-a)) - g(t)}{\varepsilon(1 - at^{-\alpha})} \\ &= D_\alpha^a(f)(t) \lim_{\varepsilon \rightarrow 0} g(t + \varepsilon t^{-\alpha}(t-a)) + f(t)D_\alpha^a(g)(t). \end{aligned}$$

Since g is continuous at t , the $\lim_{\varepsilon \rightarrow 0} g(t + \varepsilon t^{-\alpha}(t-a)) = g(t)$. This completes the proof of *iv*. The proof of the *v* is similar to *iv*. Now, we prove the result *vi*. If the function g is a constant in a neighbourhood t_0 , then $D_\alpha^a(f \circ g)(t_0) = 0$. On the other hand, we assume that g is non-constant function in the neighbourhood of t_0 . In this case, we can find an $\varepsilon_0 > 0$ such that $g(t_1) \neq g(t_2)$ for any $t_1, t_2 \in (t_0 - \varepsilon_0, t_0 + \varepsilon_0)$. Thus, since the function g is continuous at t_0 , for $t_0 > a$, $t_0^\alpha \neq a$ we get

$$\begin{aligned} D_\alpha^a(f \circ g)(t_0) &= \lim_{\varepsilon \rightarrow 0} \frac{f(g(t_0 + \varepsilon t_0^{-\alpha}(t_0-a))) - f(g(t_0))}{\varepsilon(1 - at_0^{-\alpha})} \\ &= \lim_{\varepsilon \rightarrow 0} \frac{f(g(t_0 + \varepsilon t_0^{-\alpha}(t_0-a))) - f(g(t_0))}{g(t_0 + \varepsilon t_0^{-\alpha}(t_0-a)) - g(t_0)} \cdot \frac{g(t_0 + \varepsilon t_0^{-\alpha}(t_0-a)) - g(t_0)}{\varepsilon(1 - at_0^{-\alpha})} \\ &= \lim_{\varepsilon_1 \rightarrow 0} \frac{f(g(t_0) + \varepsilon_1) - f(g(t_0))}{\varepsilon_1} \cdot \lim_{\varepsilon \rightarrow 0} \frac{g(t_0 + \varepsilon t_0^{-\alpha}(t_0-a)) - g(t_0)}{\varepsilon(1 - at_0^{-\alpha})} \\ &= f'(g(t_0)) D_\alpha^a(g)(t_0). \end{aligned}$$

This shows the Chain rule is proved.

To prove part *vii*, let $h = \varepsilon t^{-\alpha} (t - a)$ in Definition 2.1 and taking $\varepsilon = \frac{ht^\alpha}{(t-a)}$. Therefore, we get

$$\begin{aligned} D_\alpha^a (f) (t) &= \lim_{\varepsilon \rightarrow 0} \frac{f(t + \varepsilon t^{-\alpha} (t - a)) - f(t)}{\varepsilon (1 - at^{-\alpha})} \\ &= \frac{(t - a)}{(t^\alpha - a)} \lim_{\varepsilon \rightarrow 0} \frac{f(t + h) - f(t)}{h} \\ &= \frac{(t - a)}{(t^\alpha - a)} f'(t) \end{aligned}$$

since, by assumption f is differentiable at $t > 0$. This completes the proof of the theorem. □

The following theorem lists (α, a) -fractional derivative of several familiar functions.

Theorem 2.3. *Let $\alpha \in (0, 1]$, $t > a$, $t^\alpha \neq a$ and $c, n \in \mathbb{R}$. Then we have the following results*

- i.* $D_\alpha^a (t^n) = \frac{nt^{n-1}(t-a)}{(t^\alpha-a)}$
- ii.* $D_\alpha^a (1) = 0$
- iii.* $D_\alpha^a (e^{ct}) = c \frac{(t-a)}{(t^\alpha-a)} e^{ct}$
- iv.* $D_\alpha^a (\sin ct) = c \frac{(t-a)}{(t^\alpha-a)} \cos ct$
- v.* $D_\alpha^a (\cos ct) = -c \frac{(t-a)}{(t^\alpha-a)} \sin ct$
- vi.* $D_\alpha^a \left(\frac{t^\alpha}{\alpha}\right) = \frac{t^{\alpha-1}(t-a)}{(t^\alpha-a)}$.

It is easy to see from part *vi* of Theorem 2.2 that we have rather unusual results given in the following theorem.

Theorem 2.4. *Let $\alpha \in (0, 1]$ and $t > a$, $t^\alpha \neq a$. Then we have the following results*

- i.* $D_\alpha^a \left(\sin \frac{t^\alpha}{\alpha}\right) = \frac{t^{\alpha-1}(t-a)}{(t^\alpha-a)} \cos \frac{t^\alpha}{\alpha}$
- v.* $D_\alpha^a \left(\cos \frac{t^\alpha}{\alpha}\right) = -\frac{t^{\alpha-1}(t-a)}{(t^\alpha-a)} \sin \frac{t^\alpha}{\alpha}$
- vi.* $D_\alpha^a \left(e^{\frac{t^\alpha}{\alpha}}\right) = \frac{t^{\alpha-1}(t-a)}{(t^\alpha-a)} e^{\frac{t^\alpha}{\alpha}}$.

The numerical demonstrations of behaviour of a -Conformable derivative and its comparison with the Riemann derivatives for $f(t) = t^2$ and $f(t) = \sin(\pi t)$ have been presented in Figure 1-(A) and Figure 2-(A) respectively. Similarly in Figure 1-(B) and Figure 2-(B) represent the a -Conformable derivatives for different α values.

We now begin by proving the Rolle’s theorem, the mean value theorem, and the extended mean value theorem for a -conformable fractional differentiable functions.

Theorem 2.5 (Rolle’s theorem). *Let $\alpha \in (0, 1]$ and $0 \leq a < b$. If $f : [a, b] \rightarrow \mathbb{R}$ be a given function that satisfies*

- i.* f is continuous on $[a, b]$
- ii.* f is (α, a) -differentiable for some $\alpha \in (0, 1)$
- iii.* $f(a) = f(b)$.

Then , there exist $c \in (a, b)$, such that $D_\alpha^a (f) (c) = 0$.

Proof. The proof is done in a similar way in [6]. □

Theorem 2.6 (Mean Value theorem). *Let $\alpha \in (0, 1]$ and $0 \leq a < b$. If $f : [a, b] \rightarrow \mathbb{R}$ be a given function that satisfies*




FIGURE 1. (A) a -Conformable derivative, Riemann derivative and function values versus t when $a = 1$ and $\alpha = 0.25$ for $f(t) = t^2$: a -Conformable derivative (red), Riemann derivative (blue), $f(t)$ (green). (B) a -Conformable derivative of $f(t) = t^2$ for different α : $\alpha = 0.25$ (black), $\alpha = 0.50$ (blue), $\alpha = 0.75$ (red), $\alpha = 1.00$ (green).




FIGURE 2. (A) a -Conformable derivative, Riemann derivative and function values versus t when $a = 1$ and $\alpha = 0.25$ for $f(t) = \sin(\pi t)$: a -Conformable derivative (red), Riemann derivative (blue), $f(t)$ (green) (B) a -Conformable derivative of $f(t) = \sin(\pi t)$ for different α : $\alpha = 0.25$ (black), $\alpha = 0.50$ (blue), $\alpha = 0.75$ (red), $\alpha = 1.00$ (green).

- i. f is continuous on $[a, b]$
 - ii. f is (α, a) -differentiable for some $\alpha \in (0, 1)$.
- Then , there exist $c \in (a, b)$, such that

$$D_{\alpha}^a (f) (c) = \frac{f(b) - f(a) c^{\alpha-1} (c - a)}{\frac{b^{\alpha}}{\alpha} - \frac{a^{\alpha}}{\alpha}} \frac{c^{\alpha-1} (c - a)}{(c^{\alpha} - a^{\alpha})}.$$

Proof. Let's now define a new function, as follow

$$g(x) = f(x) - f(a) - \frac{f(b) - f(a)}{\frac{b^{\alpha}}{\alpha} - \frac{a^{\alpha}}{\alpha}} \left(\frac{x^{\alpha}}{\alpha} - \frac{a^{\alpha}}{\alpha} \right).$$

Since g is continuous on $[a, b]$, (α, a) -differentiable, and $g(a) = 0 = g(b)$, then by Rolle's theorem, there exist a $c \in (a, b)$ such that $D_\alpha^a(g)(c) = 0$ for some $\alpha \in (0, 1)$. Using the fact that $D_\alpha^a\left(\frac{t^\alpha}{\alpha}\right) = \frac{t^{\alpha-1}(t-a)}{(t^\alpha-a)}$, reach to the desired result. \square

Theorem 2.7 (Extended Mean Value theorem). *Let $\alpha \in (0, 1]$ and $0 \leq a < b$. If $f : [a, b] \rightarrow \mathbb{R}$ be a given function that satisfies*

- i. f, g is continuous on $[a, b]$*
 - ii. f, g is (α, a) -differentiable for some $\alpha \in (0, 1)$*
 - iii. $D_\alpha^a(g)(t) \neq 0$ for all $t \in (a, b)$.*
- Then , there exist $c \in (a, b)$ such that*

$$\frac{D_\alpha^a(f)(c)}{D_\alpha^a(g)(c)} = \frac{f(b) - f(a)}{g(b) - g(a)}.$$

Remark 2.1. *If $g(t) = \frac{t^\alpha}{\alpha}$, then this is just the statement of the Mean Value Theorem for α -conformable fractional differentiable functions.*

Proof. Let's now define a new function, as follow

$$F(x) = f(x) - f(a) - \frac{f(b) - f(a)}{g(b) - g(a)}(g(x) - g(a)).$$

Then the function F satisfies the conditions of Rolle's theorem. Thus, there exist a $c \in (a, b)$ such that $D_\alpha^a(F)(c) = 0$ for some $\alpha \in (0, 1)$. Using the linearity of D_α^a , we have

$$0 = D_\alpha^a(F)(c) = D_\alpha^a(f)(c) - \frac{f(b) - f(a)}{g(b) - g(a)}D_\alpha^a(g)(c).$$

Therefore, we get desired result. \square

3. DEFINITIONS AND PROPERTIES OF (α, a) -CONFORMABLE FRACTIONAL INTEGRAL

Now we introduce the (α, a) -conformable fractional integral (or (α, a) -fractional integral) as follows:

Definition 3.1 ((α, a) -Conformable fractional integral). *Let $\alpha \in (0, 1]$ and $0 \leq a < b$. A function $f : [a, b] \rightarrow \mathbb{R}$ is (α, a) -fractional integrable on $[a, b]$ if the integral*

$$\int_a^b f(x) d_\alpha^a x := \int_a^b f(x) \frac{(x^\alpha - a)}{x - a} dx \tag{3}$$

exists and is finite. All α -fractional integrable on $[a, b]$ is indicated by $L^1_{(\alpha, a)}([a, b])$.

Remark 3.1.

$$I^a_{(\alpha, a)}(f)(t) = \int_a^t f(x) \frac{(x^\alpha - a)}{x - a} dx,$$

where the integral is the usual Riemann improper integral, and $\alpha \in (0, 1]$.

Theorem 3.1 (Inverse property). *Let $\alpha \in (0, 1]$ and $0 \leq a < b$. Also, let $f : (a, b) \rightarrow \mathbb{R}$ be a continuous function such that $I^a_{(\alpha, a)}(f)$ exists. Then, for all $t > a$, $t^\alpha \neq a$ we have*

$$D_\alpha^a\left(I^a_{(\alpha, a)}(f)\right)(t) = f(t).$$

Proof. Since f is continuous, then $I_{(\alpha,a)}^a(f)$ is clearly differentiable. Therefore, by using *vii* of Theorem 2.2, we get

$$\begin{aligned} D_{\alpha}^a \left(I_{(\alpha,a)}^a(f) \right) (t) &= \frac{(t-a)}{(t^{\alpha}-a)} \frac{d}{dt} I_{(\alpha,a)}^a(f) (t) \\ &= \frac{(t-a)}{(t^{\alpha}-a)} \frac{d}{dt} \left(\int_a^t f(x) \frac{(x^{\alpha}-a)}{x-a} dx \right) \\ &= \frac{(t-a)}{(t^{\alpha}-a)} f(t) \frac{(t^{\alpha}-a)}{t-a} = f(t). \end{aligned}$$

This completes the proof. \square

Theorem 3.2. Let $\alpha \in (0, 1]$ and $0 \leq a < b$. Also, let $f : (a, b) \rightarrow \mathbb{R}$ be differentiable function. Then, for all $t > a$, $t^{\alpha} \neq a$ we have

$$I_{(\alpha,a)}^a(D_{\alpha}^a(f))(t) = f(t) - f(a).$$

Proof. Since f is differentiable, then, by using *vii* of Theorem 2.2, we get

$$\begin{aligned} I_{(\alpha,a)}^a(D_{\alpha}^a(f))(t) &= \int_a^t \frac{(x^{\alpha}-a)}{x-a} D_{\alpha}^a(f)(x) dx \\ &= \int_a^t \frac{(x^{\alpha}-a)}{x-a} f'(x) \frac{(x^{\alpha}-a)}{x-a} dx \\ &= f(t) - f(a). \end{aligned}$$

\square

Theorem 3.3. Let $\alpha \in (0, 1]$ and $0 \leq a < b$. Also, let $f, g : [a, b] \rightarrow \mathbb{R}$ be continuous functions. Then

- i. $\int_a^b [f(x) + g(x)] d_{\alpha}^a x = \int_a^b f(x) d_{\alpha}^a x + \int_a^b g(x) d_{\alpha}^a x$
- ii. $\int_a^b \lambda f(x) d_{\alpha}^a x = \lambda \int_a^b f(x) d_{\alpha}^a x$, $\lambda \in \mathbb{R}$
- iii. $\int_a^b f(x) d_{\alpha}^a x = - \int_b^a f(x) d_{\alpha}^a x$
- iv. $\int_a^b f(x) d_{\alpha}^a x = \int_a^c f(x) d_{\alpha}^a x + \int_c^b f(x) d_{\alpha}^a x$
- v. $\int_a^a f(x) d_{\alpha}^a x = 0$
- vi. if $f(x) \geq 0$ for all $x \in [a, b]$, then $\int_a^b f(x) d_{\alpha}^a x \geq 0$
- vii. $\left| \int_a^b f(x) d_{\alpha}^a x \right| \leq \int_a^b |f(x)| d_{\alpha}^a x$ for $x^{\alpha} > a$.

Proof. The relations follow from Definition 3.1 and Theorem 3.2, analogous properties of (α, a) -fractional integral, and the properties of section 2 for the a -conformable fractional derivative. \square

4. CONCLUDING REMARKS

In this paper a new type of fractional derivatives and integrals is proposed and tested. Relevant results such as Roll's theorem, Mean Value theorem and Extended Mean Value theorem useful for further research are also given subsequent sections. In the light of the findings above the proposed method have a number of significant implications for future practice. For instance one can present the a -conformable version of partial differential equations, Laplace transforms or related theorems.

REFERENCES

- [1] Abdeljawad, T (2015), On conformable fractional calculus, *Journal of Computational and Applied Mathematics* 279, pp. 57–66.
- [2] Anderson, D. R. (2016), Taylor’s formula and integral inequalities for conformable fractional derivatives, *Contributions in Mathematics and Engineering, in Honor of Constantin Caratheodory*, Springer, New York.
- [3] Hammad M. A. and Khalil R. (2014), Conformable fractional heat differential equations, *International Journal of Differential Equations and Applications* 13(3), pp. 177-183.
- [4] Hammad M. A. and Khalil R. (2014), Abel’s formula and wronskian for conformable fractional differential equations, *International Journal of Differential Equations and Applications* 13(3), pp. 177-183.
- [5] Iyiola O.S.and Nwaeze E.R.(2016), Some new results on the new conformable fractional calculus with application using D’Alambert approach, *Progr. Fract. Differ. Appl.*, 2(2), pp.115-122.
- [6] Khalil R., Al horani M., Yousef A. and Sababheh M.(2014), A new definition of fractional derivative, *Journal of Computational Applied Mathematics*, 264, pp. 65-70.
- [7] Katugampola U.N. (2011), New approach to a generalized fractional integral, *Appl. Math. Comput.*, 218(3), pp. 860–865.
- [8] Katugampola U.N. (2014), New approach to generalized fractional derivatives, *B. Math. Anal. App.*, 6(4), pp. 1–15.
- [9] Kilbas A. A., Srivastava H.M. and Trujillo J.J. (2016), *Theory and Applications of Fractional Differential Equations*, Elsevier B.V., Amsterdam, Netherlands.
- [10] Samko, S.G., Kilbas, A.A. and Marichev, O.I. (1993), *Fractional Integrals and Derivatives: Theory and Applications*, Yverdon: Gordon and Breach.



Mehmet Zeki SARIKAYA received his BSc (Maths), MSc (Maths) and PhD (Maths) degree from Afyon Kocatepe University, Afyonkarahisar, Turkey in 2000, 2002 and 2007 respectively. At present, he is working as a Professor in the Department of Mathematics at Duzce University (Turkey) and as a Head of Department. Moreover, he is founder and Editor-in-Chief of Konuralp Journal of Mathematics (KJM). He is the author or coauthor of more than 200 papers in the field of Theory of Inequalities, Potential Theory, Integral Equations and Transforms, Special Functions, Time-Scales.



Hüseyin BUDAK graduated from Kocaeli University, Kocaeli, Turkey in 2010. He received his M.Sc. from Kocaeli University in 2013 and PhD from Düzce University in 2017. Moreover, he works as a Assistant Professor at Düzce University. His research interests focus on functions of bounded variation, fractional calculus and theory of inequalities.



Fuat Usta received his BSc (Mathematical Engineering) degree from Istanbul Technical University, Turkey in 2009 and MSc (Mathematical Finance) from University of Birmingham, UK in 2011 and PhD (Applied Mathematics) from University of Leicester, UK in 2015. At present, he is working as a Assistant Professor in the Department of Mathematics at Düzce University (Turkey). He is interested in Approximation Theory, Multivariate approximation using Quasi Interpolation, Radial Basis Functions and Hierarchical/Wavelet Bases, High-Dimensional Approximation using Sparse Grids. Financial Mathematics, Integral Equations, Fractional Calculus, Partial Differential Equations.